

TERRO®

Outdoor Liquid Ant Baits




Directions for use

Instrucciones en Español dentro del cartón.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

- 1 Remove Stakes** by pulling them straight out from underneath cup. **DO NOT REMOVE CUP.**
- 2 Push Down** on the top of the cup, then twist cup 1/4 turn. The foil seal under the cup will be punctured and the liquid bait will drip into the reservoir in the base. **KEEP BAITS LEVEL TO AVOID SPILLING.**
- 3 Place Bait Stations** outdoors close to where ants are found entering your home, or where you see ant nests or their trails.
- 4 Secure Bait Stations** to the ground, if desired, by inserting the 2 stakes provided through the holes in the base.
- 5 Ants Will Enter** the bait station through the holes at the top of the ramps.
- 6 Monitor** regularly for activity, but do not interfere with the ants or baits. Replace baits every three months to keep ants from returning. Complete control may take up to two weeks.
- 7 Call Us** at 1-800-800-1819 if we can be of assistance.



PULL OUT STAKES



PUSH DOWN & TWIST CUP



PERIMETER DEFENSE

FOR USE IN AND AROUND: Homes, Hospitals, Restaurants, Grocery Stores, Schools, Cafeterias, Storage Areas and Food Processing Plants.

ATTRACTS AND KILLS ALL COMMON

HOUSEHOLD ANTS INCLUDING: Argentine ants, Ghost ants, Corneld ants, Pavement ants, Acrobat ants, White footed ants, Little black ants, Odorous house ants, Crazy ants, Big headed ants, and other sweet eating ants.

USING TERRO® Outdoor Liquid Ant Baits:

Since invading ants usually have outdoor nesting sites, it often helps to put the product outside, close to where ants are found entering the structure. Ants typically follow dened trails from the nest to food or water. Placing TERRO® Outdoor Liquid Ant Baits close to these trails will improve control.

CONTROLLING AND ELIMINATING ANTS:

First ants will come in large numbers, but soon disappear. This product is specially formulated to allow worker ants to consume the product and survive long enough to carry the liquid back to the nest and deliver a dose to the rest of the colony. The key to obtaining effective control is to monitor the placements and ensure there is always a fresh supply of TERRO® Outdoor Liquid Ant Baits. Complete control may take up to two weeks.

DO NOT USE in edible product areas of food handling establishments, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed. Avoid contamination of feed and foodstuffs. Place bait in areas inaccessible to children and pets.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. **STORAGE:** Store in a cool dry place. Do not store where children or animals may gain access. **PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable Container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

ACTIVE INGREDIENTS/INGREDIENTE ACTIVO:

Sodium Tetraborate Decahydrate (Borax)	5.40%
OTHER INGREDIENTS/OTROS INGREDIENTES:	94.60%
TOTAL	100.00%

KEEP OUT OF REACH
OF CHILDREN
CAUTION

MANTÉNGASE FUERA DEL
ALCANCE DE LOS NIÑOS
PRECAUCIÓN



Ant Killer

Plus Multi-Purpose Insect Control₂



DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Keep children and pets off treated area until product has been watered in and lawn is dry.

RESTRICTIONS:

- Apply this product directly to the lawn or garden area. Water treated area as directed on this label. Do not water to the point of runoff.
- Do not make applications during rain.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.
- Do not apply this product to edible growing crops or stored raw agricultural commodities used for food or feed.
- Do not allow applications to contact water inhabited by fish, such as in ornamental fish ponds that are located around structures being treated.
- Do not apply this product by aerial application.
- Use of this pesticide adjacent to water may affect aquatic organisms. To protect these organisms, do not apply this pesticide within 25 ft. of lakes, reservoirs, rivers, permanent streams, marches or natural ponds, estuaries, and commercial fish farm ponds.
- In New York State, this product may not be applied within 100 feet of a coastal marsh, or stream that drains directly into a coastal marsh.
- Observe all restrictions and precautions which appear on the label of this product.
- Do not apply more than 0.36 lbs. of the active ingredient (800 pounds of this product) per acre per year.

IMPORTANT: Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in outdoor ornamental gardens or parks, or on lawns or grounds.

GENERAL INFORMATION

For use outdoors, on lawns and as a band treatment around house foundation.

Kills ants (including Fire Ants), crickets, earwigs, fleas, roaches (German and American) and other insect pests listed in the tables below. To use this product follow these 2 easy steps – (1) Determine the size of the job and measure the area to be treated. (2) Treat as directed in rate chart. Complete coverage is important. Water in granules immediately following application.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **PESTICIDE STORAGE:** Store in a cool, dry area which is inaccessible to children and pets. Reclose container tightly after each use. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Completely empty container into application equipment. Then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **SPILLS:** If the container is leaking or material is spilled, carefully sweep up spilled product and wash exposed area. In spill or leak incidents, keep unauthorized people away.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides. **First Aid:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact Poison Control Center for emergency medical treatment at 1-800-222-1222. **IF SWALLOWED:** Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person. **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS: This product is highly toxic to fish and aquatic organisms. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind and rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

Note: Buyer assumes all responsibilities for safety and use not in strict accordance with directions.

Treatment around Foundations / Tratamiento alrededor de cimientos

Around house foundation to control pests normally found around or near foundations of houses, as well as in lawns, and certain of these pests that may enter houses.

Alrededor de los cimientos de la casa para controlar las plagas que normalmente se encuentran alrededor o cerca de los cimientos de las casas, así como en el césped y algunas de estas plagas que podrían penetrar a las casas.

Pests	Use Rate	Use Instructions
Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement, Red Imported Fire Ant, Nuisance), Fleas, Roaches (German and American), Palmetto bugs, Waterbugs, Millipedes, Mites, Silverfish, Sowbugs (Pillbugs), Crickets, Earwigs	Apply at the rate of 2 lb. to 3 lb. per 1,000 sq. ft. (an area 5' x 200') or 3.2 oz. to 4.8 oz. per 100 sq. ft. (10' x 10').	Apply a 3 to 10 foot band of granules next to the house foundation and completely distribute the granules. Water lightly immediately following application. Treat when insects first appear or when you notice lawn damage. For best results, leaf litter and debris should be removed from the foundation area. Branches and vegetation that touch the foundation should be cut back. Other potential pest harborage sites, such as deep mulch, should be raked away before application.
Plagas	Proporción de uso	Instrucciones de uso
Hormigas (argentinas, sureñas, del campo, de Allegheny, carpinteras de Florida, carpinteras negras, del maíz, de la miel, del pavimento, hormiga roja de fuego importada, molestas, pulgas, cucarachas (alemanas y americanas), de la palma, periplaneta americana, ciempiés, ácaros, lespisma, cochinilla de humedad (cochinilla), grillos, tijeretas)	Aplique a una proporción de 2 libras a 3 libras por cada 1,000 pies cuadrados (0.91 kg a 1.36 kg por 92.9 m ²) (un área de 5 pies x 200 pies) (1.5 m x 61 m) ó 3.2 onzas a 4.8 onzas por cada 100 pies cuadrados (3.2 onzas a 4.8 onzas por 9.3 m ²) (10 pies x 10 pies) (3 m x 3 m).	Aplique una franja de 3 a 10 pies (0.91 m a 3 m) de gránulos cerca del cimiento de la casa y distribuya completamente los gránulos. Riegue ligeramente de inmediato luego de la aplicación. Trate cuando los insectos aparezcan por primera vez o cuando note que hay daño en el césped. Para obtener los mejores resultados, la hojarasca y los desechos se deben retirar del área de los cimientos. Se debe cortar la vegetación y las ramas que rocen los cimientos. Antes de la aplicación, se deben barrer otros posibles lugares de anidación de plagas, como compuestos orgánicos profundos.

Spot Treatment / Tratamiento tópico

Pests	Use Rate	Use Instructions
Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement, Red Imported Fire Ant, Nuisance)	Mound Treatment: Apply 1–2 teaspoons over and around each ant hill.	Water lightly immediately following application. NOTE: Use both broadcast and spot treatment for difficult to large ant infestations. If soil is dry, water lightly before application to bring the ants closer to the surface for better control.
Fire Ants	Mound Treatment: Apply 1/2 cup over, and 2 ft. around, each fire ant mound.	Immediately following application, water in completely. Apply a minimum of 1 gallon of water to each mound area. To prevent the disturbance of ants, water gently. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. NOTE: For best results, apply in cool weather, 65°–80°F; or in early morning or late evening hours. Treat new mounds as they appear. Fast acting. Works within 24 hours.
Plagas	Proporción de uso	Instrucciones de uso
Hormigas (argentinas, sureñas, del campo, montoneras de Allegheny, carpinteras de Florida, carpinteras negras, del maíz, de la miel, del pavimento, hormiga roja de fuego importada, molesta)	Tratamiento de hormiguero: Aplique 1 a 2 cucharaditas sobre y alrededor de cada hormiguero.	Riegue ligeramente inmediatamente después de la aplicación. NOTA: Use el tratamiento por dispersión y el localizado para plagas de hormigas difíciles o grandes. Si la tierra está seca, riéguela ligeramente antes de la aplicación para acercar a las hormigas a la superficie y así obtener un mejor control.
Hormigas del fuego	Tratamiento por hormiguero: Aplique 1/2 taza sobre y 2 pies (60 cm) alrededor de cada hormiguero.	Riegue completamente inmediatamente en seguida de la aplicación. Aplique un mínimo de 1 galón de agua a cada área de hormiguero. Riegue suavemente para evitar que las hormigas se alboroten. El uso de equipo de riego con alta presión puede alborotar las hormigas y ocasionar que emigren, disminuyendo la efectividad del producto. NOTA: Para mejores resultados, aplique agua fría 65°–80° (18°–27° C) o en las horas tempranas de la mañana, o tarde en la noche. Trate los nuevos hormigueros a medida que aparezcan. Actúa rápido, en el lapso de 24 horas.

ACTIVE INGREDIENT/INGREDIENTE ACTIVO:

Lambda-cyhalothrin 0.04%

OTHER INGREDIENTS/OTROS INGREDIENTES: 99.96%

TOTAL 100.00%

KEEP OUT OF REACH
OF CHILDREN
CAUTION

MANTÉNGASE FUERA DEL
ALCANCE DE LOS NIÑOS
PRECAUCIÓN